

Chromaticity Considerations for Nonlinear Integrable Lattices

Stephen D. Webb^{#1}, David Bruhwiler¹, Alexander Valishev²,
Rami Kishek^{1,3}, Slava Danilov⁴, Sergei Nagaitsev²

¹RadiaSoft, LLC., ²FermiLab, ³University of Maryland, ⁴Oak Ridge National Lab

#swebb@radiasoft.net

18 June 2014

2014 Advanced Accelerator Concepts Workshop

Outline

- *Integrable Optics for the Intensity Frontier*
- *Conditions for integrability*
- *The Off-Momentum Problem*
- *Dispersion*
- *Chromaticity*
- *Future work*

Fermilab to Advance the Intensity Frontier

- Study of neutrino physics & rare processes
 - Requires large beam currents and high brightness (PIP-II parameters):
 - CW proton beam
 - 60-120 GeV
 - 1.2 MW
 - Intensity-dependent effects limit designs
 - beam halo
 - Laslett tune shift
- Future accelerators must mitigate these effects
 - Most collective effects arise from single betatron tunes
 - Highly nonlinear lattices will decohere collective resonances
 - Requires:
 - large tune spreads
 - bounded, (integrable?) dynamics
 - large dynamic aperture

Integrable Optics for the Intensity Frontier

- Integrable Optics Test Accelerator (IOTA)
 - Study nonlinear integrable optics of Danilov & Nagaitsev¹
 - Under construction @ FermiLab ASTA (<http://asta.fnal.gov>)
 - Implements practical lattice with
 - tune spreads of order unity
 - bounded, integrable motion
 - current plan to study single-particle dynamics
 - Other work underway (talks past and future)
 - Collective effects with intense beams (D. Bruhwiler earlier)
 - Construction of the elliptic potential magnet (F. O'Shea next)
 - Experimental program @ IOTA (S. Nagaitsev Thurs. 9:30)
 - Octupole lattices (S. Antipov Thurs. 11:20)
 - Nonlinear optics @ UMER (K. Ruisard Thurs. 11:40)

¹V. Danilov & S. Nagaitsev, “Nonlinear accelerator lattices with one and two analytic invariants”, PR ST-AB **13**, 084002 (2010).

Conditions for Integrability

- Bertrand-Darboux equation
 - Hamiltonians with 2nd invariants quadratic in momentum satisfy:

$$xy (\partial_x^2 U - \partial_y^2 U) + (y^2 - x^2 + c^2) \partial_{xy} U + 3y\partial_x U - 3x\partial_y U = 0$$

- differential equation is **linear**
 - any superposition of potentials that satisfy this differential equation will have a 2nd invariant and be integrable
- Other auxiliary conditions for accelerators:
 - matched beta functions in the drifts with these nonlinear elements
 - equal vertical and horizontal linear tunes

The Off-Momentum Problem

- Off-momentum particles couple motion to energy
 - *Linear lattice chromaticity:*
 - energy-dependent tune could cross nonlinear resonance
 - no loss of integrability (assuming linear RF bucket/coasting beam)
 - *Linear lattice dispersion:*
 - large dispersion can cause large beam size
 - *Potential problems for elliptic potential*
 - unequal tunes violates the Bertrand-Darboux equation
 - dispersion violates the equal beta function requirement

Single-turn Map





Figure from S. Nagaitsev, "IOTA Physics Goals" (2012)

Single-turn Map



Single-turn Map

- The Hamiltonian H is an invariant
 - Determines single particle dynamics
 - related to a general concept of emittance & matching
 - reduces to Courant-Snyder invariants for linear lattices
 - other invariants associated with integrable lattices
 - Contains all the information — dispersion, chromaticity...


- Normalized coordinates:

$$e^{-:H:C} = \mathcal{A} e^{-:\overline{H}:} \mathcal{A}^{-1}$$

$$e^{-:H:C} = \left(\prod_{i=0}^{N/2} \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right) \times \dots$$

$$e^{-:h_0:} \left(\prod_{i=N/2}^N \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right)_9$$

Single-turn Map



Dispersion

- After much algebra...

$$\overline{H} = \frac{\mu_0}{2} [(1 - C_x \delta) (\bar{p}_x^2 + \bar{x}^2) + (1 - C_y \delta) (\bar{p}_y^2 + \bar{y}^2) + \dots$$
$$(1 - \delta)t \int_0^{\ell_{\text{drift}}} \mathcal{V}(\bar{x} - \delta(\eta(s')/\sqrt{\beta_x(s')}), \bar{y}) ds'] + \text{h.o.t.}$$

- treating dispersion as a perturbation...

- obtain $H_0 + \delta H_1$ with H_0 integrable and

$$H_1 \approx \left(\delta(1 - \delta)t \int_0^{\ell_{\text{drift}}} \frac{\eta(s')}{\sqrt{\beta_x(s')}} ds' \right) \partial_{\bar{x}} \mathcal{V}(\bar{x}, \bar{y})$$

- Conclusions:
 - dispersion-free drifts for the magnets are ideal
 - failing that, minimize the integrated dispersion

Chromaticity

$$\overline{H} = \frac{\mu_0}{2} [(1 - C_x \delta) (\bar{p}_x^2 + \bar{x}^2) + (1 - C_y \delta) (\bar{p}_y^2 + \bar{y}^2) + \dots \\ (1 - \delta)t \int_0^{\ell_{\text{drift}}} \mathcal{V}(\bar{x} - \delta(\eta(s')/\sqrt{\beta_x(s')}), \bar{y}) ds'] + \text{h.o.t.}$$

- Bertrand-Darboux equation needs *isotropic linear term*
 - Creates non-integrable perturbation H_2
$$C_0 = \frac{1}{2}(C_x + C_y)$$
$$\Delta_C = \frac{1}{2}(C_x - C_y)$$
$$H_2 = \delta \Delta_C (\bar{p}_y^2 - \bar{p}_x^2 + \bar{y}^2 - \bar{x}^2)$$
 - Conclusions:
 - defocussing quadratic perturbation due to differing chromaticities
 - already have large tune spreads — no need to remove all the chromaticity
 - correct to make $C_x = C_y$

Conclusions

- Off-momentum dynamics breaks integrability
 - Dispersion violates the equal-beta-function requirement
 - Chromaticity introduces quadrupole perturbation which breaks the integrability
- Design to ameliorate the problem
 - Tune the chromaticities to be equal — don't have to cancel entirely!
 - conventional chromatic sextupole families suffice
 - can use weaker sextupoles, which saves more dynamic aperture
 - Use dispersion-free drifts for the nonlinear magnets
 - More study required to be sure...

Thank you for your attention

This work is supported in part by US DOE Office of Science, Office of High Energy Physics under SBIR award DE-SC0011340. We would like to thank Étienne Forest and Alex Chao for useful discussions.

18 June 2014

2014 Advanced Accelerator Concepts Workshop

Digression on Lie Operators

- Lie operators from Poisson brackets

$$\dot{z} = -\{H, z\} \mapsto \dot{z} = -:H:z \quad z(t) = e^{-:H:t} z(0)$$

- Advantages

- can multiply maps, cannot multiply Hamiltonians
- maps make coördinate transformations into similarity transformations

- Disadvantages

- a lot of formalism to get to the physics
- difficult to work with time-varying Hamiltonians

- Key Identities

- BHC Identity $e^{:C:} = e^{:A:}e^{:B:}$ $C = A + B + \frac{1}{2}[:A:B:] + \frac{1}{12}(:A:^2 B + :B:^2 A) + \dots$

- Similarity transformation $e^{:A:}e^{:B:}e^{-:A:} = \exp(:e^{:A:}B:) = e^{:B:}$

When are sextupoles optically transparent?

- Lie operator approach

$$\mathcal{M} = e^{-S_n :z^n:} e^{-:h_2:} e^{-S_n :z^n:}$$

$$e^{-:h_2:} = \mathcal{A}^{-1} \underbrace{\mathcal{R}(\theta)}_{\text{pure rotation}} \mathcal{A}$$

$$\mathcal{M} = e^{-:h_2:} \exp(-S_n :e^{:h_2:} z^n:) \exp(-S_n :z^n:)$$


$$\mathcal{M} = \mathcal{A}^{-1} \mathcal{R} \exp(-S_n : \mathcal{R}(\bar{z})^n:) \exp(-S_n : \bar{z}^n:)$$

$$\mathcal{R} \propto -1, \theta = (2n+1)\pi \implies \exp(-S_n : \mathcal{R}(\bar{z})^n:) \exp(-S_n : \bar{z}^n:) \mathcal{A} = 1$$

- Off-momentum particles do not cancel exactly because θ is energy-dependent. This is the basis of chromaticity correction.


When are sextupoles optically transparent?

- Pictorial approach (design momentum)



When are sextupoles optically transparent?

- Pictorial approach (off-momentum)



The horror...

$$\begin{aligned}\mathcal{M} &= \left(\prod_{i=0}^{N/2} \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right) e^{- : h_0 :} \left(\prod_{i=N/2}^N \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right) \\ &= \left(\prod_{i=0}^{N/2} \exp \left\{ - t_i : e^{-(i+1/2) : p^2/2 : \Delta s} \mathcal{V}_i(x, y) : \Delta s \right\} \right) \circ \\ &\quad \underbrace{e^{- : p^2/2 : \ell/2} e^{- : h_0 :} e^{- : p^2/2 : \ell/2}}_{e^{- : h_2 :}} \circ \\ &\quad \left(\prod_{i=N/2}^N \exp \left\{ - t_i : e^{(i+1/2) : p^2/2 : \Delta s} \mathcal{V}_i(x, y) : \Delta s \right\} \right)\end{aligned}$$

... the horror

$$e^{-:h_2:} = \mathcal{A} e^{-:\overline{h}_2:} \mathcal{A}^{-1}$$

Normalized coördinates

$$\mathcal{A}^{-1} = \begin{pmatrix} 1/\sqrt{\beta_x} & 0 & 0 & 0 & 0 & -\eta/\sqrt{\beta_x} \\ \alpha_x/\sqrt{\beta_x} & \sqrt{\beta_x} & 0 & 0 & 0 & -\alpha_x\eta + \beta_x\eta'/\sqrt{\beta_x} \\ 0 & 0 & 1/\sqrt{\beta_y} & 0 & 0 & 0 \\ 0 & 0 & \alpha_x/\sqrt{\beta_y} & \sqrt{\beta_y} & 0 & 0 \\ \eta' & \eta & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad \text{Courant-Snyder Parameterization}$$

$$\mathcal{M} = \left(\prod_{i=0}^{N/2} \exp \left\{ -t_i : e^{-(i+1/2) : p^2/2 : \Delta s} \mathcal{V}_i(x, y) : \Delta s \right\} \right) \circ \\ \left(\mathcal{A} e^{-:\overline{h}_2:} \mathcal{A}^{-1} \right) \circ \\ \left(\prod_{i=N/2}^N \exp \left\{ -t_i : e^{(i+1/2) : p^2/2 : \Delta s} \mathcal{V}_i(x, y) : \Delta s \right\} \right)$$

$$\begin{aligned}
& \mathcal{A}^{-1} \exp \left\{ -t_i : e^{(i+1/2) : p^2 : 2} \mathcal{V}_i(x, y) : \Delta s \right\} = \\
& \mathcal{A}^{-1} \exp \left\{ -t_i : e^{(i+1/2) : p^2 : 2} \mathcal{V}_i(x, y) : \Delta s \right\} \mathcal{A} \mathcal{A}^{-1} = \\
& \underbrace{\mathcal{A}^{-1} e^{-(i+1/2) : p^2 / 2 : \Delta s}}_{\mathcal{A}_i^{-1}} \exp \left\{ -t_i : \mathcal{V}_i(x, y) : \Delta s \right\} \underbrace{e^{(i+1/2) : p^2 / 2 : \Delta s} \mathcal{A}}_{\mathcal{A}_i}
\end{aligned}$$

The Danilov-Nagaitsy potential normalizing trick as follows:

$$A_0^{(i)} = \begin{pmatrix} 1/\sqrt{\beta_x} & 0 & 0 & 0 \\ \alpha_x/\sqrt{\beta_x} & \sqrt{\beta_x} & 0 & 0 \\ 0 & 0 & 1/\sqrt{\beta_y} & 0 \\ 0 & 0 & \alpha_x/\sqrt{\beta_y} & \sqrt{\beta_y} \end{pmatrix}$$

$$\mathcal{V}_i(x, y) = \mathcal{V}_i \left(A_0^{(i)}(x, y) \right)$$

$$\mathcal{A}^{-1} e^{-(i+1/2) : p^2/2 : \Delta s} \exp \{-t_i : \mathcal{V}_i(x, y) : \Delta s\} e^{(i+1/2) : p^2/2 : \Delta s} \mathcal{A} =$$

$$\exp \left\{ -t : \mathcal{V} \left(\bar{x} - \delta \frac{\eta}{\sqrt{\beta_x}}, \bar{y} \right) : \Delta s \right\}$$

Final transfer map in normalized coordinates

$$\left(\prod_{i=0}^{N/2} \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right) e^{- : h_0 :} \left(\prod_{i=N/2}^N \exp \left\{ - : \frac{p^2}{2} + t_i \mathcal{V}_i(x, y) : \Delta s \right\} \right) =$$

$$\mathcal{A} \exp \left\{ \sum_i -(1-\delta)t : \mathcal{V} \left(\bar{x} - \delta \frac{\eta_i}{\sqrt{\beta_i}}, \bar{y} \right) : \right\} e^{- : \bar{h}_2 :} \exp \left\{ \sum_i -(1-\delta)t : \mathcal{V} \left(\bar{x} - \delta \frac{\eta_i}{\sqrt{\beta_i}}, \bar{y} \right) : \right\} \mathcal{A}^{-1}$$

$$\bar{h}_2 = \frac{\mu_0}{2} \left[(1 - C_x \delta) (\bar{p}_x^2 + \bar{x}^2) + (1 - C_y \delta) (\bar{p}_y^2 + \bar{y}^2) \right]$$